

## LATE MIDDLE TO EARLY LATE TRIASSIC RADIOLARIAN FAUNAS FROM THE IZMIR-ANKARA SUTURE BELT IN WESTERN TURKEY: REMARKS ON THE EVOLUTION OF THE NEOTETHYAN IZMIR-ANKARA OCEAN

Uğur Kağan TEKİN<sup>1</sup> & M. Cemal GONCUOĞLU<sup>2</sup>

<sup>1</sup> Geological Engineering Department, Hacettepe University, Beytepe-Ankara, TR-06532  
[uktekin@hacettepe.edu.tr](mailto:uktekin@hacettepe.edu.tr)

<sup>2</sup> Geological Engineering Department, Middle East Technical University, Ankara, TR-06531  
[mcongcu@metu.edu.tr](mailto:mcongcu@metu.edu.tr)

Izmir-Ankara Suture Belt (IASB) in north Turkey comprises the remnants of Neotethyan Izmir-Ankara Ocean (GONCUOĞLU, 2000) and is an almost 200 km wide E-W trending belt between the Sakarya Composite Terrane in the north (GONCUOĞLU et al., 1996) and the Tauride-Anatolide Platform (TAP) in the south. Within the suture belt several tectonic units of three different main tectonic settings were recognized as thrust sheets above the units of the TAP:

1. Northern and uppermost one including more or less complete successions of the IAO lithosphere.
2. Median one with tectonic mélanges of the subduction-accretionary complex of IAO, comprising blocks of ophiolites, fore-and back-arc volcanics, radiolarian cherts, pelagic limestones and blueschists,
3. The southern and lower one belonging to the Maastrichtian –Early Paleocene flysch complex (Bornova Flysch Zone) with olistoliths and olistostromes of the former two settings and the TAP margin successions. It was formed in a peripheral foreland basin on the northern edge on the TAP, in front of the southward advancing nappes (GONCUOĞLU et al., 2006b).

The olistoliths within the melange, especially volcanics associated with sediments provide evidence for the geological evolution of the TAP continent margin as the well as the initial stages of the rifting and opening of Izmir-Ankara oceanic sea-way. Our long-lasting study along the IASB has shown that the oldest radiolarian sediments associated with volcanism are not older than Triassic. From the Bornova Flysch Zone to the NW of Manisa W Turkey) five different Triassic radiolarian assemblages have been derived from the blocks in the IASB (Tekin et al., 2006). Four of assemblages have been obtained from isolated blocks mainly composed of a radiolarian cherts- mudstones alternation. Their ages are: Late Ladinian, coinciding the *Muelleritortis cochleata* zone; early Carnian, coinciding the *Tritortis kretaensis* Zone; late Early Carnian, coinciding the “unnamed radiolarian zone” between *Tritortis kretaensis* and *Tetraporobrachia haeckeli* zones and early Middle Carnian, coinciding the lower part of *Tetraporobrachia haeckeli* zone established by KOZUR & MOSTLER (1994, 1996) and KOZUR (2003).

One continuous section (Komurcu Measured Section) with an alternation of chert and mudstone was recognized in a mega- block further NW. Diverse radiolarian faunas were obtained from this section that ranges from late Ladinian, coinciding the *Muelleritortis cochleata* zone to late Early Carnian, coinciding the “unnamed radiolarian zone” between *Tritortis kretaensis* and *Tetraporobrachia haeckeli* zones. *Pseudostylosphaera mostleri* n. sp. was defined from this section.

The new data obtained from the Bornova Flysch Zone reveals that the earliest radiolarian chert deposition to the N of the TAP commenced during the late Middle Triassic. Combined with geochemical data from the associated volcanic rocks it is shown that this event is related to the opening of the of the Izmir-Ankara Branch of Neotethys confirming our earlier suggestions (GONCUOGLU et al. 1996, 2001, 2003, 2006a; GONCUOGLU, 2000; and TEKIN et al. 2002, 2006) from the Eastern parts of IASB.

## References

- GONCUOGLU, M. C. (2000): Restoration of an alpine foreland-thrust belt: Kutahya - Bolkardag zone of the Tauride-Anatolide platform, NW Turkey.- Geology 2000, Vienna, April 14-17, 2000, Terra Nostra, Schriften der Alfred Wegener Stiftung, 2000/1: **47**, Vienna.
- GONCUOGLU, M. C., DIRIK, K. & KOZLU, H. (1996): Pre-alpine and alpine terranes in Turkey: explanatory notes to the terrane map of Turkey.- Ann. Géol. des Pays Héli., **37**: 515–536.
- GONCUOGLU, M. C., TEKIN, U. K. & TURHAN, N. (2001): Geological meaning of the late Carnian basalts blocks with radiolarites within the Late Cretaceous Central Sakarya ophiolitic complex (NW Anatolia).- Jeo 2000 Proceedings, CD-54-6: 6, Ankara.
- GONCUOGLU, M. C., TURHAN N. & TEKIN U. K. (2003): Evidence for the Triassic rifting and opening of the Neotethyan Izmir-Ankara Ocean, northern edge of the Tauride-Anatolide platform, Turkey.- Boll. della Soc. Geol. It., Spec. Vol., **2**: 203-212, Rome.
- GONCUOGLU, M. C., YALINIZ, M. K. & TEKIN, U. K. (2006a): Geochemistry, tectono-magmatic discrimination and radiolarian ages of basic extrusives within the Izmir-Ankara Suture Belt (NW Turkey): Time constraints for the Neotethyan evolution.- Ofioliti, **31**: 25-38, Firenze.
- GONCUOGLU, M. C., YALINIZ, M. K. & TEKIN, U. K. (2006b): Geochemical features and radiolarian ages of volcanic rocks from the Izmir-Ankara Suture Belt, western Turkey. - Mesozoic Ophiolite Belts of the Northern Part of the Balkan Peninsula, Proceedings: 41-44, Belgrade-Banja Luka.
- KOZUR, H. (2003): Integrated ammonoid, conodont and radiolarian zonation of the Triassic.- Hallesches Jahrb. Geowiss., **25**: 49–79, Halle.
- KOZUR, H. & MOSTLER, H. (1994): Anisian to Middle Carnian radiolarian zonation and description of some stratigraphically important radiolarians.- Geol. -Paläont. Mitt. Innsbruck, **3**: 39-255, Innsbruck.
- KOZUR, H. & MOSTLER, H. (1996): Longobardian (Late Ladinian) Oertlispongidae (Radiolaria) from the Republic of Bosnia-Herzegovina and the stratigraphic value of advanced Oertispongidae.- Geol. -Paläont. Mitt. Innsbruck, **4**: 105-193, Innsbruck.
- TEKIN, U. K., GONCUOGLU, M. C. & TURHAN, N. (2002): First evidence of Late Carnian radiolarians from the Izmir-Ankara Suture Complex, central Sakarya, Turkey: implications for the opening age of the Izmir-Ankara branch of Neo-Tethys.- Geobios, **35** (1): 127-135, Villurbanne.
- TEKIN, U. K., GONCUOGLU, M. C., OZKAN-ALTINER, S. & YALINIZ, M. K. (2006): Neotetis volkanitlerinin planktonik fosil faunasi (Radyolarya ve Planktonik Foraminifera) ile yaslandırılması, Bornova Filiş Zonu, KB Anadolu (Dating of Neotethyan volcanics by planktonic fossil faunas, Bornova Flysch Zone, NW Anatolia). TUBITAK Project No: 103Y027, Final Rep.: 1-226, Ankara (unpublished) (in Turkish with English abstract).