

TECTONIC SETTING OF SOME PALAEOZOIC METAMORPHICS IN NORTHERN ANATOLIA, TURKEY

Rojay Bora, Goncuoglu M.Cemal

Middle East Technical University,
Department of Geological Engineering,
06531 Ankara, TURKEY

ABSTRACT

In northern Anatolia along the Izmir-Ankara-Erzurum Suture, the low-grade metamorphics display strong protolithologic similarities. However, the classification of these metamorphics with respect to their protolithological stratigraphy, internal organization, age and boundary relations are still not clear due to limited number of metamorphic studies.

Amasya region in NW Central Anatolia is selected and studied to shed some light onto the metamorphic problem in northern Anatolia.

Tectonostratigraphically, three distinctly different metamorphic rock assemblages are differentiated on the basis of their internal organizations and protolithologies, as bottom to top; 1) grayish-black schists with quartz boudins and veins, 2) green schists, 3) green schists with marble blocks and/or boudins. The lower unit is originally a clastic sequence, made of shales, sandstones, cherts and calcareous clastics. It is tectonically overlain by green schists protoliths of which are volcanic and carbonate rocks. The upper unit, composed of huge marble blocks and/or lenses embedded within a volcanic sequence.

The protoliths of greenschist facies metamorphics are interpreted as a magmatic arc-related basinal being metamorphosed by regional dynamothermal metamorphism during pre-Permian. These metamorphics can protolithologically be correlated with Agvanis and Yenisehir low-grade metamorphics which all are believed to be in Carboniferous age.

INTRODUCTION

Overprinting of series of geological events caused complex terrain evolutions in northern Anatolia. As a result, it is hard and problematic to differentiate the discontinuous and scattered metamorphic units situated to the north of Izmir-Ankara-Erzincan Suture (Fig. 1).




Fig. 1- Distribution of metamorphic rocks in northern Anatolia, situated to the north of Izmir-Ankara-Erzincan Suture, irrespective to age (modified from Brinkman, 1976).

Exposures of the metamorphics are widespread in the Pontide belt (Fig. 1) (Blumenthal, 1950; Nebert, 1961; Alp, 1972; Bingol et al., 1975; Erk, 1975; Seymen, 1975; Brinkman, 1976; Ayaroglu, 1979; Kocyigit, 1979, 1987; Krushensky et al. 1980; Ozcan et al., 1980; Sengor et al., 1980; Senturk and Karakose, 1981; Tekeli, 1981; Okay, 1983, 1984, 1989; Akyurek et al.,

1984; Korkmaz and Baki, 1984; Goncuoglu et al., 1987; Yilmaz et al., 1989; Bozkurt, 1990; Okay et al., 1990; Kocyigit et al., 1991; Rojay, 1993; Genc and Yucel, 1995).

However, the differentiation of the metamorphics with respect to their protolithological stratigraphy, internal organization, age and boundary relations and tectonic settings are still not clear due to limited number of studies with well-defined petrography, geochemistry, petrofabric analyses and radiometric dating.

Various authors had proposed different age and boundary relations (Ozturk, 1979; Genc et al., 1986; Yucel et al., 1989; 1995; Genc and Yucel, 1995; Kocyigit, 1991; Kocyigit et al., 1991) different tectonic setting to the metamorphics and, therefore different scenarios to the evolution of the Tethy's in northern Anatolia (Bailey and McCallien, 1953; Bingol et al., 1973; Sengor et al., 1980, 1984, 1988; Guvenc and Konuk, 1981; Sengor and Yilmaz, 1981; Tekeli, 1981; Yilmaz et al., 1982; Sengor, 1984; Okay, 1989; Okay et al., 1990; Kocyigit et al., Genc and Yilmaz, 1995; Yucel et al., 1995).

Here, the metamorphic sequences will be stratigraphically and petrographically documented with the field data from Amasya region that may shed some light onto the evolution of metamorphics in northern Anatolia regarding to the age, boundary relations and possible tectonic setting of metamorphics.




Fig. 2- Measured reference section of the low-grade metamorphics in Amasya (Locality: southern slope of Karasanlar ridge, E of Amasya).

The metamorphics which are one of the basement units of the Mesozoic sequences in Pontides consist dominantly of metavolcanic, metapyroclastic and meta-volcaniclastic rocks, metaclastic rocks and metacarbonates-meta-cherts with a minimum observable thickness of about 310 m.

Tectonostratigraphically, three distinctly different metamorphic rock assemblages are differentiated on the basis of their internal organizations and protolithologies. These subunits depending upon their protolithologies are: (1) Grayish-black schists with quartz boudins and veins, (2) Green schists, (3) Green schists with marble blocks and or boudins (Fig. 2).

In the measured section, the sequence starts with the grayish black and intensely deformed schists which are included in muscovite-chlorite-calcite-quartz-albite schist having zircon, tourmaline and biotite clasts experienced a schistose texture (meta-clastic), muscovite bearing quartz schist, muscovite±albite±chlorite-quartz schist with schistose texture (meta-chert), quartz-albite-muscovite schist with schistose texture (meta-pelitic rock) and tourmaline bearing chlorite-sericite-albite calcschist with slaty cleavage (meta-carbonate siltstone) paragenesis. This facies is characterized by the presence of slaty structure and quartz (silica) boudins (Fig. 2).

Tectonically, the sequence continues with light green-green foliated slates (chlorite-actinolite-epidote-muscovite-albite schist with schistose texture) with a metatuffaceous unit (stilpnomelane-chlorite-epidote-albite schist with schistose texture) and is followed by dominantly green-yellowish green-white massive to laminated marble-slate alternation (epidote-actinolite-chlorite schist with nematoblastic chlorite and Na-amphibole). Green, thin-bedded to laminated siliceous marble and marble alternating with green slates (epidote-actinolite-chlorite schist, quartz-muscovite-chlorite ± apatite schist and chlorite-quartz-albite-calcite ± epidote ± opaque schist with schistose texture) grades upward into green foliated schists (epidote-actinolite-chlorite schist with nematoblastic texture) and to thin laminated, banded, mylonitized marble bands (chlorite-albite-calcite ± pyrite clac-schist with schistose structure) alternating with green slates (epidote-actinolite-chlorite ± albite schist with schistose texture).

The top of the sequence consists dominantly of green-light green thin bedded-laminated schists (chlorite-epidote-actinolite-albite ± leucoxene schist rich in Na-amphiboles with nematoblastic texture, epidote-chlorite-Na-amphibole an epidote-actinolite-chlorite schists with nematoblastic texture) with dismembered grayish white-yellow marble blocks and/or boudins (Fig. 2).

In other sections, the following paragenesis are obtained green biotite bearing chlorite-actinolite-calcite-albite ± epidote ± sphene schist with schistose structure (meta-tuff), green biotite-bearing epidote-actinolite-albite-sphene schist with lepidoblastic texture (meta-diabase), chlorite-actinolite-albite ± epidote ± sphene schist with porphyroblastic texture (meta-tuff), epidote-actinolite-albite ± muscovite ± sphene schist with lepidoblastic texture (meta-gabbro), epidote-chlorite-actinolite-albite ± sphene schist with lepidoblastic texture (meta-diabase), actinolite-albite-epidote ± biotite ± sphene (few) schist with lepidoblastic texture (meta-"gabbroic" rock), epidote-actinolite-albite ± green biotite ± sphene ± chlorite schist with schistose texture (meta-volcanic rock), actinolite - epidote-albite ± sphene ± green biotite ± chlorite schist with relict volcanic texture (albite-epidote filled amygdaloids) (meta-pillow basalt). In addition to above mentioned protoliths, green meta-cherts (some are radiolarian cherts with poorly preserved radiolaria tests) are significant facies of the metabasics.

The schistose texture is observed in meta-tuffs and meta-sedimentary rocks, lepidoblastic texture in meta-diabasic-gabbroic rocks, porphyroblastic texture in meta-tuffs with nematoblastic chlorite and Na-amphibole. The bedding attitudes and schistosity planes have conformable attitudes and the pillow structures of the basalts are well preserved as well. However, cataclastic and mylonitic textures are also well developed at the contacts of various different meta-facies.

CORRELATION, AGE RELATIONS AND INTERPRETATION

Some metamorphics are correlative units in the northern Anatolia especially the Agvanis metamorphics (Okay, 1984; Rojay, 1985), Tokat metamorphics (Blumenthal, 1950; Alp, 1972; Ozcan et al., 1980; Rojay, 1993) and Yenisehir metamorphics (Genc et al., 1986; Yilmaz et al., 1989; Kocyigit et al., 1991) on the basis of degree and type of metamorphism, protolithostratigraphy, tectonic and stratigraphic setting. However, the controversial, confusive and misuse of the nomenclature, especially the "Karakaya" terminology, cease most of the studies from a regionwide correlation.

The widely excepted age for the low-grade metamorphics in northern Anatolia is pre-Liassic due to regionwide Liassic discordance.

However, the equivalent low-grade metamorphics in the northwestern Anatolia are unconformably overlain by Triassic clastics (Krushensky et al., 1980; Gence et al., 1986; Kocyigit et al., 1991) and by Permian clastics and carbonates (Genc et al., 1986; Goncuoglu et al., 1987). Studies carried out on the fragments of Triassic clastics showed that Carboniferous and Permian age fragments are present besides the metamorphic quartz and rock fragments (Kocyigit et al., 1991; Rojay, 1993).

An age of 272 ± 3 Ma is obtained by K/Ar radiometric dating from the granitoids having cross-cutting relationship with low-grade metamorphics (Cogulu and Krummenacher, 1967). Therefore the age of the metamorphism should be older than age of the granitic intrusions which is early Permian.

Collectively, the metamorphics should be older than Permian, based on the overlying Permian sedimentary units and cross-cutting relationship with granitoids. However, the regional considerations and correlations with protolithologically similar, fossiliferous equivalent units in Kutahya-Bolkardagi Belt (Goncuoglu et al., 1992; Goncuoglu, this volume) indicate that the deposition and metamorphism of the assemblage will be contemporaneous with Carboniferous.

Depending on the protolithology of the sequence, various facies of the metamorphics were deposited in a basin where intense subalkaline to alkaline basaltic volcanism was active. The extensive distribution of mafic volcanics with pillow basalts and possible radiolarites (pelagic influx) may manifest a deep sea depositional setting in an active continental marginal basin (Dickinson and Selley, 1981). Therefore these metamorphics can be interpreted as a magmatic arc related basinal sequences which were resulted from a regional dynamothermal metamorphism in green schist facies.

REFERENCES

- AKYUREK, B., BILGINER, E., AKBAS, B., HEPSEN, N., PEHLIVAN, S., SUNU, O., SOYSAL, Y., DAGER, Z., CATAL, E., SOZERI, B., YILDIRIM, H. & HAKYEMEZ, Y., (1984). Ankara-Elmadag-Kalecik dolayinin temel jeolojik ozellikleri. *Jeoloji Muhendisligi*, 20, 31-46.
- ALP, D., (1972). Amasya yoresinin jeolojisi, *Ist Univ.Fen Fak. Monogr.*, 22, 101.
- ALTINLI, I.E., (1975). Bilecik Jurasigi, Cumhuriyetin 50. Yili Yerbilimleri Kongresi, M.T.A. Enst. Ankara, 103-111.
- ALTINLI, I.E., GURPINAR, O. & ERISEN, S. (1970). Geology of the Erenkoy-Deresakari area (Bilecik), *Ist Univ. Fen Fak Mec.*, 35, 1, 77-83.
- AYAROGLU, H., (1979). Bozuyuk metamorfiteilerinin (Bilecik) petrokimyasal ozellikleri. *T.J.K. Bult.*, 22, 1, 101-108.
- BAILEY, E.B. & Mc CALLIAN, W.J., (1953). Serpentine Lavas, the Ankara Melange and the Anatolian Thrust. *Phil. Trans R. Soc. London*, no. LXII, pp. 404-442.
- BINGOL, E., AKYUREK, B. & KORKMAZER, B., (1975). Biga Yarimadasinin jeolojisi ve Karakaya formasyonunun bazi ozellikleri. Cumhuriyetin 50. Yili Yerbilimleri Kongresi, M.T.A. Ozel Baski, 70-76, Ankara.
- BLUMENTHAL, M., (1950). Orta ve Asagi Yesilirmak bolgesinin (Tokat-Amasya-Havza-Erbaa-Niksar)

- jeolojisi hakkında. M.T.A. Yayini, D/4, 153 p.
- BOZKURT, E., (1990). Geology of the Almus Fault Zone (AFZ) (Almus-Tokta). M.S. Thesis, M.E.T.U., Ankara, 118 p.
- BRINKMANN, R., (1972). Mesozoic troughs and crustal structure in Anatolia. *Geol. Soc. Am. Bull.*, 83, 819-826.
- BRINKMANN, R., (1976). Geology of Turkey. Elsevier Scientific Publishing Company, Amsterdam, 153p.
- COGULU, E. & KRUMMENACKER, DD., (1967). Problemes geochronometriques dans la partie NW de l'Anatolie Centrale (Turquie) *Bull. Suisse de Mineral. Petrographie.*, 47, 2, 825-831.
- DICKINSON, W.R., & SELLY, D.R., (1979). Structure and stratigraphy of forearc regions. *Bull. Am. Assoc. of Pet. Geol.*, 63, 1, 2-31.
- ERK, S., (1975). Ankara yoresinin genc Paleozoyikin stratigrafisi. TUBITAK 5. Bilim Kong. Teblig., Izmir, Yerbilimleri Seksiyonu, 1-29.
- GENC, C. & YUCEL. Y., (1995). Evolution of the Triassic continental margin of NW Anatolia. *Tectonophysics*, 243, 155-171.
- GENC, S., SELCUK, H., CEVHER, F., GO-ZLER, Z., KARAMAN, T., BILGI, C., VE AKGOREN, F., (1986). Inegol (Bursa)-Pazaryeri (Bilecik) arasinin jeolojisi. M.T.A. Rap. 255, 68 p.
- GONCUOGLU, E., ETENDIL, M., TEKELI, O., AKSAY, A., KUSCU, I. & URGUN, B.M. (1987). Geology of the Armutlu peninsula. Guide book for the field excursion along Western Anatolia, Turkey, IGCP Project 5, 12-18.
- GONCUOGLU, M.C., OZCAN, A., TURHAN, N. & ISIK, A., (1992). Stratigraphy of the Kuta-hya Region. ISGB. International Symposium on the Geology of the Black Sea Region. A geotraverse across Tethyan Suture Zones in NW Anatolia. 3-8.
- GONCUOGLU, M.C. (this volume) Rock Units and Geodynamic Evolution of the Central Sakarya Terrane and its Correlation with the Balkan Terranes. Carpatho-Balkan Meeting, 1995.
- GUVENÇ, T. & KONUK, Y.T., (1981). Triyas boyunca tortullasma ve kiriklanma, Ege-Anadolu Kirik Zonu. H.U. Yerbilimleri, 7, 43-53.
- KOCYIGIT, A., (1979). Tekneli Bolgesinin (Tokat guneyi) tektonik Ozelligi. TUBITAK Proje Raporu, No. TBAG-262, 63 p.
- KOCYIGIT, A., (1987). Hasanoglan (Ankara) yoresinin tektonostratigrafisi: Karakaya orojenik kusaginin evrimi. H.U. Yerbilimleri, 14, 269-293.
- KOCYIGIT, A., (1991). First Remark on the Geology of Karakaya Basin. Karakaya Orogen and pre-Jurassic nappes in eastern Pontides, Turkey. *Geologica Romana*, 27, 3-11.
- KOCYIGIT, A., KAYMAKCI, N., ROJAY, B.F., OZCAN, E., DIRIK, K., & OZCELIK, Y., (1990). Inegol-Bilecik-Bozuyuk arasinda kalan alanin jeolojik etudu. M.E.T.U.-T.P.A.O. Re-search Project, No. 90-03-09-01-05.
- KORKMAZ, S. & BAKI, Z., (1984). Demirozu (Bayburt) guneyinin stratigrafisi. *T.J.K. Bult.* 5, 107-115.
- KRUHENSKY, R.D., AKCAY, Y., & KARAEĞE, E., (1980). Geology of the Karalar-Yesiller area, north west Anatolia, Turkey. *U.S. Geol. Surv. Bull.*, 1462, 72.
- NEBERT, K., (1961). Kelkit cayi ve Kizilirmak (Kuzeydogu Anadolu) nehirleri mecras bolgelerin jeolojik yapisi. M.T.A. Derg., 57, 1-49.
- OKAY, A.I., (1983). Agvanis metamorfikleri ve cevre kayalarinin jeolojisi M.T.A. Derg., 99/100, 51-71.
- OKAY, A.I., (1984). Kuzeybati Anadolu'da yer alan metamorfik kusaklar. Ketin Simpozyumu, T.J.K. Ozel Baskisi, 83-92.
- OKAY, A.I., (1989). Tectonic units and sutures in the Pontides, northern Turkey. In A.M.C. SENGOR (ed.), Tectonic Evolution of the Tethyan Region, 109-115.
- OKAY, A.I., SIYAKO, M., & BURKAN, K.A., (1990). Biga yarimadasinin jeolojisi ve tektonik evrimi. T.P.J.D. Bult., 2, 1, 83-121.
- OZCAN, A., ERKAN, A., KESKIN, A., KESKIN, E., ORAL, A. OZER, S., SUMEGEN, M., TE-KELI, O., (1980). Amasya-Turhal arasindaki bolgenin jeolojisi. M.T.A. Rap. No: 6722.
- OZTURK, A., (1979). Ladik-Destek yoresinin stratigrafisi. *T.J.K. Bult.*, 22, 27-34.
- ROJAY, B., (1985). The tectonostratigraphic characteristics of the Kelkit valley subzone of the NAF. Ms. Thesis, METU, Ankara, 105.
- ROJAY, B., (1993). Tectonostratigraphy and neotectonic characteristics of the southern margin of Merzifon-Suluova Basin (Central Pontides, Amasya). PhD. Thesis, METU, Ankara, 215.
- SEYMEN, I., (1975). Kelkit vadisi kesiminde Kuzey Anadolu Fay Zonunun tektonik ozelligi. Istanbul Tek. Univ. Maden Fak. yayini, 192.
- SENGOR, A.M.C., (1984), The Cimmeride Orogenic System and the tectonics of Eurasia. *Geol. Soc. Am. Spec. Paper* 195, 82.

- SENGOR, A.M.C., YILMAZ, Y., & KETIN, I., (1980). Remnants of a pre-Late Jurassic ocean in northern Turkey: fragments of Permian-Triassic Paleo-Tethys?. *Geol. Soc. Am. Bull.*, 91, 599-609.
- SENGOR, A.M.C. & YILMAZ, Y. (1981). Tethyan evolution of Turkey: a plate tectonic approach. *Tectonophysics*, 75, 181-241.
- SENGOR, A.M.C., YILMAZ, Y., & SUNGURLU, O., (1984). Tectonics of the Mediterranean Cimmerides: nature and evolution of the western termination of Paleo-Tethys. In J.E. DIXON & A.H.F. ROBERTSON (eds.), The geological evolution of the Eastern Mediterranean, *Geol. Soc. London Spec. Publ.* 17, 77-112.
- SENGOR, A.M.C., ALTINER, D., CIN, A., USTAOMER, T., & HSU, K.J., (1988). Origin and assembly of the Tethyside orogenic collage at the expense of Gondwana Land. In AUD-LEY-CHARLES, M.G. and HALLAM, A. (eds.), Gondwana and Tethys, *Geol. Soc. London Spec. Publ.* 37, 119-181.
- SENTURK, K. & KARAKOSE, C., (1981). Orta Sakarya bölgesinde Liyas öncesi ofiyolitlerin ve mavi sistlerin oluşumu ve yerleşmesi. *T.J.K. Bult.*, 24, 1-10.
- TEKELI, O., (1981), Subduction complex of pre-Jurassic age, north Anatolia, Turkey. *Geology*, 9, 68-72.
- YILMAZ, Y., (1981). Sakarya kitasi güney kenarının tektonik evrimi. *Ist. Univ. Yerbilimleri*, 1, 1-2, 33-52.
- YILMAZ, Y., TUYSUZ, O., GOZUBOL, A.M. & YIGITBAS, E., (1981). Abant (Bolu)-Dokurcun (Sakarya) arasında Kuzey Anadolu Fay Zonunun kuzey ve güneyinde kalan tektonik birliklerin jeolojik evrimi. *Istanbul Yerbilimleri, I.U.Y.B.F. Yayın organı*, 2, 3-4, 239-261.
- YILMAZ, Y., GURPINAR, O., GENÇ, C., AÖZCU, M., YILMAZ, K., SEKER, H., YIGITBAS, E., & KESKİN, M., (1989). Armutlu yarımadası ve dolayının jeolojisi. I.T.U.-T.P.A.O. pro-jesi. 2796, 210.
- YILMAZ, Y., GENÇ, C., YIGITBAS, E., BOZCU, M., & YILMAZ, K., (1995). Geological evolution of the late Mesozoic continental margin of NW Anatolia. *Tectonophysics*, 243, 155 -171.