

# Fixed Point Free Action on Groups of Odd Order

DEDICATED TO THE MEMORY OF BRIAN HARTLEY

by

Gülin Ercan and İsmail Ş. Güloğlu

## Introduction

Let  $G$  be a finite solvable group and  $A$  be a finite group acting fixed point freely on  $G$ . A longstanding conjecture is that if  $(|G|, |A|) = 1$ , then the Fitting length of  $G$  is bounded by the length  $\ell(A)$  of the longest chain of subgroups of  $A$ . By an elegant result due to Bell and Hartley [1], it is known that any finite nonnilpotent group  $A$  can act fixed point freely on a solvable group  $G$  of arbitrarily large Fitting length with  $(|G|, |A|) \neq 1$ . We expect that the conjecture is true when the coprimeness condition is replaced by the assumption that  $A$  is nilpotent. This question is still unsettled except for cyclic groups  $A$  of order  $pq$  and  $pqr$  for pairwise distinct primes  $p, q$  and  $r$  ([3], [5]).

In the present paper we establish the conjecture without the coprimeness condition when  $A$  is a finite abelian group of square free odd exponent not divisible by 3 and  $|G|$  is odd. This improves the bound given in Theorem 3.4 of [7]. We also improve some bounds given in Theorems 8.4, 8.5 of [2].

Namely, we shall prove the following:

**Theorem A.** Let  $G$  be a finite group of odd order and  $A$  be a finite abelian group of square free odd exponent not divisible by 3. If  $C_G(A) = 1$ , then  $f(G) \leq \ell(A)$ .

**Theorem B.** Let  $H$  be a group of odd order not divisible by 3. Suppose that its Carter subgroups have a normal complement  $G$ . If  $C$  is a Carter subgroup of  $H$ , then  $f(G) \leq 2(2^{\ell(C)} - 1)$ .

**Theorem C.** Let  $G$  be a group of odd order not divisible by 3. If  $C$  is a Carter subgroup of  $G$ , then  $f(G) \leq 4(2^{\ell(C)} - 1) - \ell(C)$ .

Except for the following, the notation and terminology are as in [2].

For any group  $G$ ,  $\tilde{G}$  denotes the Frattini factor group of  $G$ .

Let  $K$  be a group acting on finite solvable groups  $H$  and  $G$ . We say  $(K \text{ on } G)$  and  $(K \text{ on } H)$  are weakly equivalent if each nontrivial irreducible section of  $(K \text{ on } G)$  is  $K$ -isomorphic to an irreducible section of  $(K \text{ on } H)$  and vice versa. We write  $(K \text{ on } H) \equiv_w (K \text{ on } G)$  if  $(K \text{ on } H)$  is weakly equivalent to  $(K \text{ on } G)$ .

### Some Remarks

Let  $K, L, G$  and  $H$  be groups.

(a) If  $(K \text{ on } G) \equiv_w (K \text{ on } H)$ , then  $(L \text{ on } G) \equiv_w (L \text{ on } H)$  for each  $L \leq K$ .

(b) Let  $L$  act on  $K$  and  $K$  act on  $G$  and  $H$ . If  $(K \text{ on } G) \equiv_w (K \text{ on } H)$ , then  $(K \text{ on } G)^\ell \equiv_w (K \text{ on } H)^\ell$  for each  $\ell \in L$ .

(c) Let  $V$  be a completely reducible  $kG$ -module for a field  $k$  and let  $L$  act on  $G$ . Let  $\ell \in L$  and  $V_\ell$  denote the  $kG$ -module with respect to  $(G \text{ on } V)^\ell$ . Assume that  $(G \text{ on } V) \equiv_w (G \text{ on } V)^\ell$ . Let  $M \leq G$  such that  $M$  is  $\langle \ell \rangle$ -invariant, and  $W$  be the sum of all irreducible  $kG$ -submodules of  $V$  on which  $M$  acts nontrivially. Then  $W = W^\# = W_\ell$  as subspaces where  $W^\#$  stands for the sum of all irreducible  $kG$ -submodules of  $V_\ell$  on which  $M$  acts nontrivially.

Note that  $W$  and  $W_\ell$  need not be isomorphic as  $kG$ -modules.

**Lemma 1:** Let  $S \langle \alpha \rangle$  be a group where  $S \triangleleft S \langle \alpha \rangle$ ,  $S$  is an  $s$ -group for some prime  $s$ ,  $\Phi(S) \leq Z(S)$ ,  $\langle \alpha \rangle$  is cyclic of order  $p$  for an odd prime  $p$ . Suppose that  $V$  is a  $kS \langle \alpha \rangle$ -module for a field  $k$  of characteristic different from  $s$ . Then  $C_V(\alpha) \neq 0$  if one of the following is satisfied:

- (i)  $[Z(S), \alpha]$  is nontrivial on  $V$ .
- (ii)  $[S, \alpha]^{p-1}$  is nontrivial on  $V$  and  $p = s$ .

Furthermore, if  $S \langle \alpha \rangle$  acts irreducibly on  $V$  or the characteristic of  $k$  is different from  $p$ , then we also have  $(\mathcal{C} \text{ on } C_V(\alpha)) \equiv_w (\mathcal{C} \text{ on } V)$  where  $\mathcal{C} = C_D(\alpha)$

$$\text{for } D = \begin{cases} S & \text{when (i) holds} \\ [S, \alpha]^{p-1} & \text{when (ii) holds.} \end{cases}$$

**Proof:** ([2], Proposition 3.10)

**Lemma 2:** (Lemma 5.30 in [2]). Let  $S \triangleleft S \langle \alpha \rangle$  where  $\langle \alpha \rangle$  is cyclic of prime order and let  $V$  be an irreducible  $kS \langle \alpha \rangle$ -module. If  $E$  is an  $\langle \alpha \rangle$ -invariant subgroup of  $Z(S)$  and  $U$  is a nonzero  $E \langle \alpha \rangle$ -submodule of  $V$ , then  $\text{Ker}(E \text{ on } V) = \text{Ker}(E \text{ on } U)$ .

**Lemma 3:** Let  $S \langle \alpha \rangle$  be a group such that  $S \triangleleft S \langle \alpha \rangle$  where  $\langle \alpha \rangle$  is of prime order  $p$ . Suppose that  $V$  is a  $kS \langle \alpha \rangle$ -module for a field  $k$  of characteristic different from  $p$ , and  $\Omega$  is an  $S \langle \alpha \rangle$ -stable subset of  $V^*$ . Set  $V_0 = \cap \{\text{Ker } f \mid f \in \Omega - C_\Omega(\alpha)\}$ . If there exists a nonzero  $f$  in  $\Omega$  and  $x \in S$  such that  $f(V_0) \neq 0$  and  $[x, a, \alpha] \notin C_S(f)$  for each  $1 \neq a \in \langle \alpha \rangle$ , then  $C_V(\alpha) \not\subseteq V_0$ .

**Proof:** Since  $f(V_0) \neq 0$ , it follows that  $f \in C_\Omega(\alpha)$  and so  $C_S(f)$  is normalized by  $\langle \alpha \rangle$ . The assumption  $[x, a, \alpha] \notin C_S(f)$  for each  $1 \neq a \in \langle \alpha \rangle$  yields that  $[x, a] \notin C_S(f)$  for each  $1 \neq a \in \langle \alpha \rangle$ . Then  $bx f \notin C_\Omega(\alpha)$  for each  $b \in \langle \alpha \rangle$ . Set  $g = \sum_{b \in \langle \alpha \rangle} bx f$ . It is clear that  $g \in C_\Omega(\alpha)$  and so  $[V, \alpha] \subseteq \text{Ker } g$ . Since  $V = [V, \alpha] \oplus C_V(\alpha)$ , either  $g = 0$  or  $C_V(\alpha) \not\subseteq \text{Ker } g$ . If the latter holds, then  $C_V(\alpha) \not\subseteq V_0$  as claimed, because  $V_0 \subseteq \text{Ker}(bx f)$  for each  $b \in \langle \alpha \rangle$ . Hence we may assume that  $g = 0$ . Now  $0 = x^{-1}g = f + \sum_{1 \neq b \in \langle \alpha \rangle} [x, b]f$  and then  $f = - \sum_{1 \neq b \in \langle \alpha \rangle} [x, b]f$ . Since  $[x, b, \alpha] \notin C_S(f)$  by the hypothesis, we have  $[x, b]f \notin C_\Omega(\alpha)$  for each  $1 \neq b \in \langle \alpha \rangle$ . Then  $f(V_0) = 0$ . This contradiction completes the proof.  $\square$

The following result is a generalization of Theorem 2.1.A in [6].

**Theorem 1:** Let  $S \langle \alpha \rangle$  be a group such that  $S \triangleleft S \langle \alpha \rangle$ ,  $S$  is an  $s$ -group,  $\langle \alpha \rangle$  is cyclic of order  $p$  for odd primes  $s$  and  $p$  with  $p \geq 5$ ,  $\phi(\phi(S)) = 1$ ,  $\phi(S) \leq Z(S)$ .

Suppose that  $k$  is a field of characteristic not dividing  $ps$  and  $V$  is a  $kS \langle \alpha \rangle$ -module such that  $[S, \alpha]^{p-1}$  acts nontrivially on each irreducible submodule of  $V|_S$ .

Let  $\Omega$  be an  $S < \alpha >$ -stable subset of  $V^*$  which linearly spans  $V^*$  and set  $V_0 = \cap \{ \text{Ker} f \mid f \in \Omega - C_\Omega(\alpha) \}$ . Then  $C_V(\alpha) \not\subseteq V_0$  and  $(C_D(\alpha) \text{ on } C_V(\alpha)/C_{V_0}(\alpha)) \equiv_w (C_D(\alpha) \text{ on } V)$  where  $D = \begin{cases} [S, \alpha]^{p-1} & \text{when } s = p \\ S & \text{otherwise.} \end{cases}$

**Proof:** Assume that the theorem is false and consider a counterexample with  $\dim V + |S < \alpha >|$  minimal. Set  $X = C_V(\alpha)/C_{V_0}(\alpha)$  and  $\mathcal{C} = C_D(\alpha)$ .

*Claim 1.* We may assume that  $S$  acts faithfully and  $S < \alpha >$  acts irreducibly on  $V$  and  $k$  is a splitting field for all subgroups of  $S < \alpha >$ .

Put  $\bar{S} = S/\text{Ker}(S \text{ on } V)$ . By induction applied to the action of  $\bar{S} < \alpha >$  on  $V$ , we get  $C_V(\alpha) \not\subseteq V_0$  and  $(C_{\bar{D}}(\alpha) \text{ on } X) \equiv_w (C_{\bar{D}}(\alpha) \text{ on } V)$ . As  $\bar{\mathcal{C}} = \overline{C_D(\alpha)} \leq C_{\bar{D}}(\alpha)$ , we have obtained  $(\mathcal{C} \text{ on } X) \equiv_w (\mathcal{C} \text{ on } V)$ . Thus we may assume that  $S$  is faithful on  $V$ .

Since  $V$  is completely reducible as an  $S < \alpha >$ -module, we have a collection  $\{V_1, \dots, V_\ell\}$  of irreducible  $S < \alpha >$ -submodules of  $V$  such that  $V = \bigoplus_{i=1}^{\ell} V_i$ . Now  $[S, \alpha]^{p-1}$  acts nontrivially on each irreducible constituent of  $V_i|_S$  and hence  $[S, \alpha]^{p-1}$  acts nontrivially on each  $V_i$  for  $i = 1, \dots, \ell$ . It is easy to observe that  $\Omega|_{V_i}$  is an  $S < \alpha >$ -stable subset of  $V_i^*$  and  $\langle \Omega|_{V_i} \rangle = V_i^*$  for each  $i = 1, \dots, \ell$ . If  $V$  is not irreducible as an  $S < \alpha >$ -module, we apply induction to the action of  $S < \alpha >$  on  $V_i$  for each  $i$  and get  $C_{V_i}(\alpha) \not\subseteq (V_i)_0$  and  $(\mathcal{C} \text{ on } C_V(\alpha)/C_{(V_i)_0}(\alpha)) \equiv_w (\mathcal{C} \text{ on } V_i)$ . Set  $X_i = C_{V_i}(\alpha)/C_{V_i \cap V_0}(\alpha)$ . Now  $(\mathcal{C} \text{ on } X_i) \equiv_w (\mathcal{C} \text{ on } V_i)$  since  $(V_i)_0 = \cap \{ \text{Ker} g \mid g \in \Omega_i - C_{\Omega_i}(\alpha) \} \supseteq V_i \cap V_0$ . As  $V = \bigoplus_{i=1}^{\ell} V_i$  and  $X \cong \bigoplus_{i=1}^{\ell} X_i$ , it follows that  $(\mathcal{C} \text{ on } X) \equiv_w (\mathcal{C} \text{ on } V)$ . Therefore we can regard  $V$  as an irreducible  $S < \alpha >$ -module.

*Claim 2.*  $[Z(S), \alpha, \alpha] = 1$ .

Assume the contrary. Set  $S_1 = Z(S)\mathcal{C}$ . Then  $S_1$  is an  $\langle \alpha \rangle$ -invariant subgroup of  $S$  and  $V|_{S_1 \langle \alpha \rangle}$  is completely reducible. Note that  $\mathcal{C} \triangleleft S_1 \langle \alpha \rangle$ . Let  $V_i$  be an irreducible  $S_1 \langle \alpha \rangle$ -submodule of  $V$  and  $W$  be a homogeneous component of  $V_i|_{\mathcal{C}}$ .

Now  $Z(S) \langle \alpha \rangle \leq C_{S_1 \langle \alpha \rangle}(\mathcal{C}) \leq N_{S_1 \langle \alpha \rangle}(W)$ . This yields that  $V_i|_{\mathcal{C}}$  is homogeneous. We also observe that  $\text{Ker}(Z(S) \text{ on } V_i) = \text{Ker}(Z(S) \text{ on } V) = 1$  by applying Lemma 2 to the action of  $S < \alpha >$  on  $V$ .

Since  $[Z(S), \alpha] \neq 1$ ,  $[Z(S_1), \alpha]$  is nontrivial on  $V_i$ . Applying Lemma 1 to the action

of  $S_1 \langle \alpha \rangle$  on  $V_i$ , we obtain  $C_{V_i}(\alpha) \neq 0$ . If  $C_{V_i}(\alpha) \not\subseteq V_0$ , it follows that  $(\mathcal{C} \text{ on } C_{V_i}(\alpha)/C_{V_i \cap V_0}(\alpha)) \equiv_w (\mathcal{C} \text{ on } V_i)$  as  $V_i|_{\mathcal{C}}$  is homogeneous. This forces that there is an irreducible  $S_1 \langle \alpha \rangle$ -submodule  $V_i$  of the completely reducible module  $V|_{S_1 \langle \alpha \rangle}$  such that  $C_{V_i}(\alpha) \subseteq V_0$ . Since  $0 \neq C_{V_i}(\alpha)$ , we have  $V_i \cap V_0 \neq 0$ . Set  $\Omega_i = \Omega|_{V_i}$ . Now  $\Omega_i$  is an  $S_1 \langle \alpha \rangle$ -stable subset of  $V_i^*$ , and  $(V_i)_0 = \cap \{\text{Ker } h | h \in \Omega_i - C_{\Omega_i}(\alpha)\} \neq 0$  as  $V_i \cap V_0 \subseteq (V_i)_0$ . Let  $f \in \Omega$  be such that  $f((V_i)_0) \neq 0$ . Then  $f_i = f|_{V_i} \in C_{\Omega_i}(\alpha)$ . Consider  $\langle f_i \rangle = \{cf_i | c \in k\}$ , a  $C_{Z(S)}(f_i) \langle \alpha \rangle$ -submodule of  $V_i^*$ . Appealing to Lemma 2 together with  $\langle f_i \rangle$  and  $C_{Z(S)}(f_i)$ , we get  $C_{Z(S)}(f_i) = \text{Ker}(C_{Z(S)}(f_i) \text{ on } V_i^*) = 1$ . On the other hand, there exists  $x \in Z(S)$  such that  $[x, \alpha, \alpha] \neq 1$ , as  $[Z(S), \alpha, \alpha] \neq 1$ . It follows that  $[x, a, \alpha] \neq 1$  for any  $1 \neq a \in \langle \alpha \rangle$ , that is  $[x, a, \alpha] \notin C_{S_1}(f_i)$ , for any  $1 \neq a \in \langle \alpha \rangle$ . Now Lemma 3 applied to the action of  $S_1 \langle \alpha \rangle$  on  $V_i$ , together with  $f_i$  and  $\Omega_i$ , gives that  $C_{V_i}(\alpha) \not\subseteq (V_i)_0$ . This is a contradiction as  $V_i \cap V_0 \subseteq (V_i)_0$  and  $C_{V_i}(\alpha) \subseteq V_0$ . Thus we have the claim.

*Claim 3.  $s \neq p$ .*

Assume that  $s = p$ . Since  $[S, \alpha]^{p-1} \neq 1$ ,  $[S, \alpha]^{p-3} \neq 1$ . Set  $S_1 = [S, \alpha]^{p-3}$ . We can prove that  $[S_1, [S, \alpha]^{p-1}] \leq [\phi(S), \alpha]^{p-3} = 1$  (see ([2] 5.37)). Hence  $[S, \alpha]^{p-1} \leq Z(S_1)$ .

We have a collection  $\{V_1, \dots, V_\ell\}$  of irreducible  $S_1 \langle \alpha \rangle$ -modules such that  $V = \bigoplus_{i=1}^{\ell} V_i$ . Fix  $i \in \{1, \dots, \ell\}$ . We notice that  $\mathcal{C} = C_{[S, \alpha]^{p-1}}(\alpha) \triangleleft S_1 \langle \alpha \rangle$  implying  $V|_{\mathcal{C}}$  is completely reducible. In particular,  $\mathcal{C} \leq Z(S_1 \langle \alpha \rangle)$  and so  $V_i|_{\mathcal{C}}$  is homogeneous.

Set  $X_i = C_{V_i}(\alpha)/C_{V_i \cap V_0}(\alpha)$  and assume that  $(\mathcal{C} \text{ on } X_i) \not\equiv_w (\mathcal{C} \text{ on } C_{V_i}(\alpha))$ . If  $[S, \alpha]^{p-1}$  is trivial on  $V_i$ , then  $\mathcal{C}$  acts trivially on  $V_i$ , and this contradicts the assumption. Hence  $[S, \alpha]^{p-1}$  is not trivial on  $V_i$ . If  $V_i \cap V_0 = 0$ , then  $(\mathcal{C} \text{ on } X_i) \equiv_w (\mathcal{C} \text{ on } C_{V_i}(\alpha))$ , and again we have a contradiction. Hence,  $V_i \cap V_0 \neq 0$ , and there exists some  $f \in \Omega$  such that  $f(V_i \cap V_0) \neq 0$ . Now  $f \in C_{\Omega}(\alpha)$ . Set  $f|_{V_i} = f_i$ . Now  $\langle f_i \rangle = \{cf_i | c \in k\}$  is a  $C_{[S, \alpha]^{p-1}}(f_i) \langle \alpha \rangle$ -submodule of  $V_i^*$ . Appealing to Lemma 2, we get  $C_{Z(S_1)}(f_i) = \text{Ker}(C_{Z(S_1)}(f_i) \text{ on } V_i^*)$ . We also have  $C_{[S, \alpha]^{p-1}}(f_i) \leq C_{Z(S_1)}(f_i)$ . Thus  $C_{[S, \alpha]^{p-1}}(f_i)$  is properly contained in  $[S, \alpha]^{p-1}$ , that is, there is  $1 \neq y \in [S, \alpha]^{p-1} - C_{[S, \alpha]^{p-1}}(f_i)$ , and  $x \in [S, \alpha]^{p-3}$  such that  $y = [x, \alpha, \alpha]$ . It follows that  $1 \neq [x, a, \alpha] \notin C_{[S, \alpha]^{p-1}}(f_i)$  for any  $1 \neq a \in \langle \alpha \rangle$ . Now we can apply Lemma 3 to the

action of  $S_1 \langle \alpha \rangle$  on  $V_i$  together with  $\Omega_i = \Omega|_{V_i}$  and  $f_i$ , and obtain that  $C_{V_i}(\alpha) \not\subseteq V_0$ . As  $V_i|_{\mathcal{C}}$  is homogeneous, we already have  $(\mathcal{C} \text{ on } X_i) \equiv_w (\mathcal{C} \text{ on } C_{V_i}(\alpha))$ .

Therefore we conclude that  $(\mathcal{C} \text{ on } C_V(\alpha)/C_{V_0}(\alpha)) \equiv_w (\mathcal{C} \text{ on } C_V(\alpha))$ . Appealing to Lemma 1 together with  $V$  and  $S \langle \alpha \rangle$ , we also see that  $C_V(\alpha) \neq 0$  and  $(\mathcal{C} \text{ on } C_V(\alpha)) \equiv_w (\mathcal{C} \text{ on } V)$  hold. Thus  $(\mathcal{C} \text{ on } V) \equiv_w (\mathcal{C} \text{ on } C_V(\alpha)/C_{V_0}(\alpha))$ . Since  $[S, \alpha]^{p-1} \neq 1$  and  $s = p$ ,  $\mathcal{C} \neq 1$ . Hence  $\mathcal{C}$  is nontrivial on  $V$  and so is on  $C_V(\alpha)/C_{V_0}(\alpha)$ . This supplies  $C_V(\alpha) \not\subseteq V_0$ , a contradiction

*Claim 4. The theorem follows.*

Now  $s \neq p$  and  $[\phi(S), \alpha] = 1$ . Then  $\phi(S) \leq Z(S \langle \alpha \rangle)$  and so  $S$  is a central product of  $[S, \alpha]$  and  $C_S(\alpha)$ . As  $\mathcal{C} = C_S(\alpha) \triangleleft S \langle \alpha \rangle$ ,  $V|_{\mathcal{C}}$  is completely reducible. In fact,  $V|_{\mathcal{C}}$  is homogeneous, because any homogeneous component is stabilized by  $S \langle \alpha \rangle$  as  $\mathcal{C}$  is centralized by  $[S, \alpha] \langle \alpha \rangle$ . It follows that  $(\mathcal{C} \text{ on } C_V(\alpha)/C_{V_0}(\alpha)) \equiv_w (\mathcal{C} \text{ on } V)$  if  $C_V(\alpha) \not\subseteq V_0$  holds. Hence  $C_V(\alpha) \subseteq V_0$ . Note that  $C_V(\alpha) \neq 0$ , because otherwise we would have obtained  $s = 2$  as  $[S, \alpha]$  is nontrivial on  $V$ . Then there exists  $0 \neq f \in C_{\Omega}(\alpha)$  with  $f(V_0) \neq 0$ . Now  $C_{Z(S)}(f) = \text{Ker}(C_{Z(S)}(f) \text{ on } V^*) = 1$  by Lemma 2. It follows that  $C_{Z([S, \alpha])}(f) = 1$ , as  $[C_S(\alpha), [S, \alpha]] = 1$ . Then  $C_{[S, \alpha]}(f)$  is properly contained in  $[S, \alpha]$ . Let  $M$  be a maximal  $\alpha$ -invariant subgroup of  $[S, \alpha]$  containing  $C_{[S, \alpha]}(f)$ . The abelian group  $[S, \alpha]/M = \overline{[S, \alpha]}$  forms an irreducible  $\langle \alpha \rangle$ -module on which  $\langle \alpha \rangle$  acts fixed point freely. Thus we have  $[\bar{x}, a] \neq 0$  for any  $0 \neq \bar{x} \in \overline{[S, \alpha]}$ . It follows that  $[\bar{x}, a, \alpha] \neq 0$  for each  $1 \neq a \in \langle \alpha \rangle$ . Put  $\bar{x} = xM$  for  $x \in [S, \alpha]$ . Then  $[x, a, \alpha] \notin M$ . In particular,  $[x, a, \alpha] \notin C_{[S, \alpha]}(f)$  for each  $1 \neq a \in \langle \alpha \rangle$ . Recall that  $V|_{\mathcal{C}}$  is homogeneous. Then Lemma 3 applied to the action of  $S \langle \alpha \rangle$  on  $V$  gives that  $C_V(\alpha) \not\subseteq V_0$ . This contradiction completes the proof of Theorem 1.  $\square$

**Theorem 2:** Let  $S \langle \alpha \rangle$  be a group such that  $S \triangleleft S \langle \alpha \rangle$ ,  $S$  is an  $s$ -group,  $\langle \alpha \rangle$  is cyclic of order  $p$  for distinct primes  $s$  and  $p$ ,  $\phi(\phi(S)) = 1$ ,  $\phi(S) \leq Z(S)$ . Suppose that  $V$  is an irreducible  $kS \langle \alpha \rangle$ -module on which  $[S, \alpha]$  acts nontrivially where  $k$  is a field of characteristic different from  $s$ . Then

$$[V, \alpha]^{p-1} \neq 0 \quad \text{and} \quad (C_S(\alpha) \text{ on } V) \equiv_w (C_S(\alpha) \text{ on } [V, \alpha]^{p-1})$$

unless  $p$  is a Fermat prime,  $s = 2$  and  $[\tilde{S}, \alpha]$  is an irreducible  $\langle \alpha \rangle$ -module.

**Proof:** ([2], Proposition 3.10)

Now we are ready to prove our key result, which improves Theorem 3.1 in [6] obtained by pursuing the idea in Dade's work [2].

**Theorem 3:** Let  $G \triangleleft GA$  and  $\langle z \rangle \trianglelefteq A$  of prime order  $p$  with  $p \geq 5$ . Suppose that  $P_1, \dots, P_t$  is an  $A$ -Fitting chain of  $G$  such that  $[P_1, z] \neq 1$ ,  $P_i$  is a  $p_i$ -group where  $p_i$  is an odd prime for each  $i = 1, \dots, t$ , and  $t \geq 3$ . Then there are sections  $D_{i_0}, \dots, D_t$  of  $P_{i_0}, \dots, P_t$ , respectively, forming an  $A$ -Fitting chain of  $G$  such that  $z$  centralizes each  $D_j$  for  $j = i_0, \dots, t$  where  $i_0 = \begin{cases} 2 & \text{if } p_1 \neq p \\ 3 & \text{if } p_1 = p \end{cases}$ .

**Proof:** Let  $p_{t+1}$  be a prime different from  $p_t$  and let  $P_{t+1}$  stand for the regular  $\mathbb{Z}_q[P_t P_{t+1} A]$ -module. We shall add  $P_{t+1}$  to the given chain and define subspaces  $E_i$  of  $P_i$  for each  $i = 1, \dots, t+1$  as follows:  $E_1 = P_1$ ,  $E_i = [\mathcal{X}_i, E_{i-1}]$  for  $i = 2, \dots, t+1$ , where  $\mathcal{X}_i/\phi(P_i)$  is the sum of all ample<sup>1</sup> irreducible  $E_{i-1} \langle z \rangle$ -submodules of  $\tilde{P}_i$ : It is easy to observe that for each  $i = 2, \dots, t+1$ ,  $E_i$  are all  $E_{i-1}A$ -invariant subgroups of  $P_i$  and  $\tilde{E}_i$  is a direct sum of ample irreducible  $E_{i-1} \langle z \rangle$ -submodules.

We now define subgroups  $F_i$  of  $E_i$  for  $i = 1, \dots, t+1$  as follows:

$$\begin{aligned} F_1 &= \{1\} \\ F_i &= C_{E_i}(z) \quad \text{if } p_i \neq p \quad \text{and } i \geq 2 \\ F_2 &= C_{[E_2, z]^{p-1}}(z) \quad \text{if } p_2 = p \\ F_i &= [[E_i, z]^{p-1}, F_{i-1}] \quad \text{if } p_i = p \quad \text{and } i \geq 3 \end{aligned}$$

It can also be easily seen that for each  $i = 2, \dots, t+1$ ,  $F_i$  is  $F_{i-1}A$ -invariant and is centralized by  $z$ .

We next define the sections  $D_i$  by  $D_i = F_i/\text{Ker}(F_i \text{ on } \tilde{E}_{i+1})$  for  $i = 2, \dots, t$  and claim that they form an  $A$ -chain each of its sections is centralized by  $z$ , as desired.

---

<sup>1</sup>Let  $V$  be an irreducible  $G \langle \alpha \rangle$ -module where  $G \triangleleft G \langle \alpha \rangle$  and  $\langle \alpha \rangle$  is cyclic of prime order  $p$ . We say  $V$  is an ample  $G \langle \alpha \rangle$ -module if  $[G, \alpha]^{p-1}$  acts nontrivially on  $V$ . Notice that when  $|G|$  is odd, this coincides with the definition of an ample module given in [2].

We proceed from this point by assuming that we can prove the following two claims whose proofs will follow later.

**Claim 1:** Assume that  $i \geq 2$  and  $p_i \neq p$ . If  $E_i \neq 1$ , then  $D_i$  is a nontrivial  $F_{i-1}$ -invariant section such that  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } \tilde{D}_i)$

**Claim 2:** Assume that  $i \geq 2$  and  $p_i = p$ . If either  $i = 2$  or  $D_{i-1} \neq 1$ , then  $\text{Ker}(F_i \text{ on } \tilde{E}_{i+1}) = 1$ ,  $D_i = F_i \neq 1$  and  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } \tilde{F}_i)$ .

We first prove the theorem in the case  $p_1 \neq p$ .

Now  $E_1 = P_1$  and  $[E_1, z]^{p-1} = [E_1, z] \neq 1$ . Then the faithful action of  $P_1$  on  $\tilde{P}_2 = [\tilde{P}_2, [E_1, z]] \oplus C_{\tilde{P}_2}([E_1, z])$  forces that  $\tilde{E}_2 \neq 0$ , that is,  $\tilde{P}_2$  contains an irreducible ample  $E_1 \langle z \rangle$ -submodule. If  $p_2 \neq p$ , we apply Claim 1 to the action of  $E_1 \langle z \rangle$  on  $\tilde{E}_2$  and obtain that  $D_2$  is a nontrivial section of  $E_2$ . If  $p_2 = p$ , we also have  $D_2 = F_2 \neq 1$  by Claim 2. Thus we have seen that  $D_2 \neq 1$  in any case.

Suppose that  $D_{i-1} \neq 1$  for some  $i \geq 3$ . Then  $E_i \neq 1$ . Appealing again to Claim 1 and Claim 2, respectively, when  $p_i \neq p$  and  $p_i = p$ , we see that  $D_i$  is a nontrivial  $F_{i-1}$ -invariant section and  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } \tilde{D}_i)$  for each  $i \geq 2$ . It follows that  $D_{i-1} = F_{i-1}/\text{Ker}(F_{i-1} \text{ on } \tilde{D}_i)$  normalizes  $D_i = F_i/\text{Ker}(F_i \text{ on } \tilde{E}_{i+1})$  and  $\text{Ker}(D_{i-1} \text{ on } D_i) = 1$  for each  $i = 3, \dots, t$ .

We also have  $\phi(D_i) \leq Z(D_i)$ ,  $\phi(\phi(D_i)) = 1$  and  $[\phi(D_i), D_{i-1}] = 1$  for  $i = 2, \dots, t$ .

It remains to prove that  $(D_{i-1} \text{ on } \tilde{D}_i)$  is weakly  $D_{i-2}$ -invariant for  $i = 4, \dots, t$ . Since  $(P_{i-1} \text{ on } \tilde{P}_i)$  is weakly  $P_{i-2}$ -invariant,  $(E_{i-1} \text{ on } \tilde{P}_i)$  is weakly  $F_{i-2}$ -invariant by Remark (a), that is,  $(E_{i-1} \text{ on } \tilde{P}_i) \equiv_w (E_{i-1} \text{ on } \tilde{P}_i)^x$  for each  $x \in F_{i-2}$ . Then  $\mathcal{X}_i/\phi(P_i) = (\mathcal{X}_i/\phi(P_i))_x$  by Remark (c) and so  $(E_{i-1} \text{ on } \tilde{E}_i) \equiv_w (E_{i-1} \text{ on } \tilde{E}_i)^x$ . Hence  $(E_{i-1} \text{ on } \tilde{E}_i)$  is weakly  $F_{i-2}$ -invariant. This gives that  $(F_{i-1} \text{ on } \tilde{E}_i)$  is weakly  $F_{i-2}$ -invariant, too. As  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } \tilde{D}_i)$  holds, it also follows that  $(F_{i-1} \text{ on } \tilde{D}_i)$  is weakly  $F_{i-2}$ -invariant by Remark (b). Consequently we have obtained that  $(D_{i-1} \text{ on } \tilde{D}_i)$  is weakly  $D_{i-2}$ -invariant, proving the theorem when  $p_1 \neq p$ .

Finally we assume that  $p_1 = p$ , and consider the chain  $P_2, \dots, P_t$ . Note that  $[P_2, z] \neq 1$ , because otherwise  $[P_1, z] = 1$  by the three subgroup lemma. Since  $p_2 \neq p$ , the above argument gives an  $A$ -Fitting chain  $D_3, \dots, D_t$  whose terms are all centralized



by  $z$ . This completes the proof of Theorem 3.  $\square$ .

*Proof of Claim 1.*

We have  $E_{i-1} \neq 1$  as  $[E_i, E_{i-1}] = E_i$ . Also  $\text{Ker}(E_i \text{ on } \mathcal{X}_{i+1}/\phi(P_{i+1})) = \text{Ker}(E_i \text{ on } E_{i+1}) = \text{Ker}(E_i \text{ on } \tilde{E}_{i+1})$ . Appealing to Remark (c) together with  $V = \tilde{P}_{i+1}$ ,  $G = P_i$ ,  $L = F_{i-1}$  and  $M = [E_i, z]$ , we see that  $\text{Ker}(E_i \text{ on } \tilde{E}_{i+1})$  is  $F_{i-1}$ -invariant. This yields that  $D_i = F_i/\text{Ker}(F_i \text{ on } \tilde{E}_{i+1})$  is  $F_{i-1}$ -invariant, as  $F_{i-1}$  normalizes  $F_i$ .

We know that  $\tilde{E}_i = \bigoplus_{j=1}^{\ell} W_{i_j}$  where  $W_{i_1}, \dots, W_{i_\ell}$  are irreducible ample  $E_{i-1} < z >$ -submodules. Set  $W_{i_j} = U_j/\phi(E_i)$  for each  $j = 1, \dots, \ell$ . Since  $\tilde{P}_{i+1}|_{E_i}$  is completely reducible and  $E_i$  is faithful on  $\tilde{P}_{i+1}$ , there exists at least one irreducible component of  $\tilde{P}_{i+1}|_{E_i}$  on which  $U_j$  acts nontrivially. Let  $\mathfrak{N}_j$  denote the set of all such components of  $\tilde{P}_{i+1}|_{E_i}$ .

There are two cases:

Either (I) there is at least one  $N$  in  $\mathfrak{N}_j$  on which  $\phi(E_i)$  acts trivially,

or

(II) there is no  $N$  in  $\mathfrak{N}_j$  on which  $\phi(E_i)$  acts trivially.

In the latter case, a closer look at the members of  $\mathfrak{N}_j$  gives the following:

*Let  $N$  be an irreducible component of  $\tilde{E}_{i+1}|_{E_i}$ . Then  $N \in \mathfrak{N}_j$  iff  $\phi(E_i)$  acts nontrivially on  $N$ .*

This is an immediate consequence of a more general fact stated as follows:

*Lemma.* Assume  $p_i \neq p$  and let  $W$  be an irreducible submodule of  $\tilde{P}_{i+1}|_{E_i}$ . If  $\phi(E_i)$  acts nontrivially on  $W$ , then so does  $[E_i, z]$ .

To prove this lemma, let  $W$  be an irreducible submodule of  $\tilde{P}_{i+1}|_{E_i}$  on which  $\phi(E_i)$  acts nontrivially and  $[E_i, z]$  acts trivially. Then there exists an  $E_i A$ -submodule  $X$  of  $\tilde{P}_{i+1}$  such that  $W$  is isomorphic to an irreducible  $E_i$ -submodule of  $X$ . Since  $X|_{E_i}$  is completely reducible, there is a collection  $\{U_1, \dots, U_s\}$  of homogeneous  $E_i$ -modules such that  $X = \bigoplus_{i=1}^s U_i$ . Assume that  $U_1$  is a sum of isomorphic copies of  $W$ . Then  $\text{Ker}(E_i \text{ on } X) = \bigcap_{a \in A} \text{Ker}(E_i \text{ on } U_1)^a = \bigcap_{a \in A} \text{Ker}(E_i \text{ on } W)^a$ .

Put  $K = \text{Ker}(\phi(E_i) \text{ on } X)$ .  $K$  is an  $A$ -invariant normal subgroup of  $E_i$ . Furthermore,  $K$  is  $E_{i-1}$ -invariant because  $[\phi(E_i), E_{i-1}] = 1$ . Set  $\bar{E}_i = E_i/K$  and

$\overline{\overline{E}}_i = \overline{E}_i/\text{Ker}(\overline{E}_i \text{ on } X)$ . Note that  $E'_i = \phi(E_i)$  since  $C_{E_i/E'_i}(E_{i-1}) = 0$ . Now  $\overline{E}_i$  is nonabelian, because otherwise  $E'_i = \phi(E_i) = K$ , which is not the case. It follows that  $V = \overline{E}_i/Z(\overline{E}_i) \neq 0$ . Obviously we have  $Z(\overline{\overline{E}}_i) \subseteq Z(\overline{E}_i)$ . On the other hand if  $Z(\overline{\overline{E}}_i) = \overline{\overline{C}} = \overline{C}/\text{Ker}(\overline{C} \text{ on } X)$ , then  $[\overline{C}, \overline{E}_i] \leq \text{Ker}(\overline{E}_i \text{ on } X) \cap \phi(\overline{E}_i) = 1$ , because  $\phi(\overline{E}_i) = \phi(E_i/K)$  is faithful on  $X$ . Therefore  $\overline{C} \leq Z(\overline{E}_i)$ , that is,  $Z(\overline{\overline{E}}_i) = \overline{Z(\overline{E}_i)}$ .

Also note that  $\text{Ker}(\overline{E}_i \text{ on } X) \subset Z(\overline{E}_i)$ : Because otherwise there is  $\overline{x} \in \text{Ker}(\overline{E}_i \text{ on } X) - Z(\overline{E}_i)$  and so there is  $\overline{y} \in \overline{E}_i$  such that  $1 \neq [\overline{x}, \overline{y}]$ . Now  $[\overline{x}, \overline{y}]$  is a nontrivial element of  $\phi(\overline{E}_i)$  acting trivially on  $X$ . This contradicts the fact that  $\phi(\overline{E}_i)$  is faithful on  $X$ .

Thus  $Z(\overline{\overline{E}}_i) = Z(\overline{E}_i)/\text{Ker}(\overline{E}_i \text{ on } X)$ . We conclude that  $\overline{E}_i/Z(\overline{E}_i)$  and  $\overline{\overline{E}}_i/Z(\overline{\overline{E}}_i)$  are  $\langle z \rangle$ -isomorphic modules. Since  $\langle z \rangle$  is trivial on  $\overline{\overline{E}}_i$ , it is trivial on  $V$  also. An application of the three subgroup lemma supplies that  $[E_{i-1}, z]$  is also trivial on  $V$ . It follows that  $[E_{i-1}, z]$  is trivial on each of the  $E_{i-1} \langle z \rangle$ -composition factors of  $V$ . Note that  $V$  is a nonzero quotient module of  $\tilde{E}_i$ . Since  $\tilde{E}_i$  is a direct sum of ample irreducible  $E_{i-1} \langle z \rangle$ -submodules, so is  $V$ , that is,  $[E_{i-1}, z]^{p-1}$  and hence  $[E_{i-1}, z]$  is nontrivial on  $V$ , a contradiction completing the proof of Lemma.

Now we can proceed with the proof. Recall that we are studying the case (II), that is,  $\phi(E_i)$  is nontrivial on each member of  $\mathfrak{N}_j$ . Thus  $U_j$  is trivial on each irreducible component  $N$  of  $\tilde{P}_{i+1}|_{E_i}$  lying outside  $\tilde{E}_{i+1}$ , because otherwise  $N \in \mathfrak{N}_j$  implying that  $\phi(E_i)$  and hence  $[E_i, z]$  is nontrivial on  $N$ , a contradiction. It follows that

$$1 = \text{Ker}(U_j \text{ on } \tilde{P}_{i+1}) = \text{Ker}(U_j \text{ on } \tilde{E}_{i+1}) \text{ when (II) holds.}$$

Now suppose that  $\text{Ker}(U_j \text{ on } \tilde{E}_{i+1}) = 1$  for each  $j = 1, \dots, s$  and  $\text{Ker}(U_j \text{ on } \tilde{E}_{i+1}) \neq 1$  for each  $j = s+1, \dots, \ell$ .

For each  $j = s+1, \dots, \ell$ , set  $\Omega_j = \{f \in W_{i_j}^* \mid \text{There exists } N \text{ in } \mathfrak{N}_j \text{ on which } \phi(E_i) \text{ acts trivially and } \text{Ker}(U_j \text{ on } N)/\phi(E_i) \subseteq \text{Ker} f\}$ . Now for each  $N$  in  $\mathfrak{N}_j$  on which  $\phi(E_i)$  acts trivially,  $\text{Ker}(U_j \text{ on } N)/\phi(E_i)$  is proper in  $W_{i_j}$  and hence is contained in a maximal subspace  $M$ . Therefore  $\Omega_j \neq \{0\}$ . Also  $\Omega_j$  is  $E_{i-1} \langle z \rangle$ -invariant. This yields that  $\langle \Omega_j \rangle = W_{i_j}^*$ , by the irreducibility of  $W_{i_j}^*$  as an  $E_{i-1} \langle z \rangle$ -module.

Now for each  $j = 1, \dots, \ell$ , we set  $K_j = \text{Ker}(U_j \text{ on } \tilde{E}_{i+1})$ . Then  $K_j \phi(E_i)/\phi(E_i) \subseteq$

$(W_{i_j})_0$ : If not, then

$j \in \{s+1, \dots, \ell\}$  and there exist  $x \in K_j$ ,  $f \in \Omega_j - C_{\Omega_j}(z)$  such that  $f(x\phi(E_i)) \neq 0$ . By the definition of  $\Omega_j$ , we can find an irreducible submodule  $N$  of  $\tilde{P}_{i+1}|_{E_i}$  on which  $U_j$  is nontrivial,  $\phi(E_i)$  is trivial and  $\text{Ker}(U_j \text{ on } N)/\phi(E_i) \subseteq \text{Ker} f$ . Then  $x \notin \text{Ker}(U_j \text{ on } N)$ . As  $x \in \text{Ker}(U_j \text{ on } \tilde{E}_{i+1})$ ,  $N$  lies outside  $\tilde{E}_{i+1}|_{E_i}$ , that is,  $[E_i, z]^{p-1} = [E_i, z]$  acts trivially on  $N$ . Thus  $[U_j, z]$  is trivial on  $N$  and so  $f \in C_{\Omega_j}(z)$ , a contradiction.

Since  $W_{i_j}$  is an irreducible  $E_{i-1} < z >$ -module,  $W_{i_j}|_{E_{i-1}}$  decomposes into a direct sum of homogeneous  $E_{i-1}$ -modules which are permuted transitively by  $< z >$ . Since  $[E_{i-1}, z]^{p-1}$  is nontrivial on at least one of these components, it is nontrivial on all of them. It follows that  $[E_{i-1}, z]^{p-1}$  acts nontrivially on each irreducible component of  $W_{i_j}|_{E_{i-1}}$  for each  $j = 1, \dots, \ell$ .

Let  $\Omega_j$  denote the whole of  $W_{i_j}^*$  when  $j \in \{1, \dots, s\}$ . Appealing to Theorem 1 for each  $j = 1, \dots, \ell$  together with the action of

$E_{i-1} < z >$  on  $W_{i_j}$  and the corresponding  $\Omega_j$ , we see that  $C_{W_{i_j}}(z) \not\subseteq (W_{i_j})_0$  and  $(F_{i-1} \text{ on } C_{W_{i_j}}(z)/C_{(W_{i_j})_0}(z)) \equiv_w (F_{i-1} \text{ on } W_{i_j})$ .

We shall now observe that for each  $j = 1, \dots, \ell$ ,  $(F_{i-1} \text{ on } W_{i_j}) \equiv_w (F_{i-1} \text{ on } C_{W_{i_j}}(z))$ : If  $p_{i-1} = p$  or  $[Z(E_{i-1}), z]$  is nontrivial on  $W_{i_j}$ , this holds by Lemma 1. Assume that  $p_{i-1} \neq p$  and  $[Z(E_{i-1}), z] \leq K = \text{Ker}(E_{i-1} \text{ on } W_{i_j})$ . Since  $[E_{i-1}, z]$  is nontrivial on  $W_{i_j}$  and  $p_{i-1}$  is odd, it can be easily seen that  $C_{W_{i_j}}(z) \neq 0$ . Put  $\bar{E}_{i-1} = E_{i-1}/K$ . As  $\overline{\phi(E_{i-1})} = \phi(\bar{E}_{i-1}) \leq Z(\bar{E}_{i-1} < z >)$ ,  $\bar{E}_{i-1}$  is a central product of  $[\bar{E}_{i-1}, z] < z >$  and  $C_{\bar{E}_{i-1}}(z)$ . Then  $C_{\bar{E}_{i-1}}(z) \triangleleft \bar{E}_{i-1} < z >$  and  $W_{i_j}|_{C_{\bar{E}_{i-1}}(z)}$  is homogeneous. We have  $\bar{F}_{i-1} \leq C_{\bar{E}_{i-1}}(z)$  yielding that  $(\bar{F}_{i-1} \text{ on } W_{i_j}) \equiv_w (\bar{F}_{i-1} \text{ on } C_{W_{i_j}}(z))$ . Thus  $(F_{i-1} \text{ on } W_{i_j}) \equiv_w (F_{i-1} \text{ on } C_{W_{i_j}}(z))$ .

Now  $(F_{i-1} \text{ on } C_{W_{i_j}}(z)) \equiv_w (F_{i-1} \text{ on } C_{W_{i_j}}(z)/C_{(W_{i_j})_0}(z)) \equiv_w (F_{i-1} \text{ on } W_{i_j})$  holds, for each  $j = 1, \dots, \ell$ .

Set  $L_j = \text{Ker}(C_{U_j}(z) \text{ on } \tilde{E}_{i+1})$ . Notice that any nontrivial irreducible  $F_{i-1}$ -submodule of  $C_{W_{i_j}}(z)/C_{(W_{i_j})_0}(z)$  is  $F_{i-1}$ -isomorphic to an irreducible  $F_{i-1}$ -submodule of  $C_{U_j}(z)/L_j$ . Therefore any nontrivial irreducible  $F_{i-1}$ -submodule of  $W_{i_j}$  is  $F_{i-1}$ -isomorphic to an

irreducible  $F_{i-1}$ -submodule of  $C_{U_j}(z)/L_j$ . On the other hand any nontrivial irreducible  $F_{i-1}$ -submodule of  $C_{U_j}(z)/L_j$  is  $F_{i-1}$ -isomorphic to an irreducible  $F_{i-1}$ -submodule of  $C_{U_j}(z)$  and hence to an irreducible  $F_{i-1}$ -submodule of  $W_{i_j}$ . This shows that  $(F_{i-1} \text{ on } W_{i_j}) \equiv_w (F_{i-1} \text{ on } C_{U_j}(z)/L_j)$  for each  $j = 1, \dots, \ell$ .

As  $\tilde{E}_i = \bigoplus_{j=1}^{\ell} W_{i_j}$  and  $C_{\tilde{E}_i}(z) = \bigoplus_{j=1}^{\ell} C_{W_{i_j}}(z) = \bigoplus_{j=1}^{\ell} C_{U_j}(z)\phi(E_i)/\phi(E_i)$ , we have  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } C_{E_i}(z)/\text{Ker}(C_{E_i}(z) \text{ on } \tilde{E}_{i+1}))$ . Notice that  $D_i = C_{E_i}(z)/\text{Ker}(C_{E_i}(z) \text{ on } \tilde{E}_{i+1})$ . Hence  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } D_i) \equiv_w (F_{i-1} \text{ on } \tilde{D}_i)$ , because  $[\phi(D_i), F_{i-1}] = 1$ . Since  $C_{W_{i_j}}(z) \not\subseteq (W_{i_j})_0$  we have  $F_i = C_{E_i}(z) \not\subseteq \text{Ker}(E_i \text{ on } \tilde{E}_{i+1})$  and so  $D_i \neq 1$ , completing the proof of Claim 1.  $\square$

*Proof of Claim 2.*

Suppose that  $p_i = p$  for some  $i \geq 2$ . If  $i \neq 2$ , assume that  $D_{i-1} \neq 1$ . Now  $\text{Ker}([E_i, z]^{p-1} \text{ on } \tilde{E}_{i+1}) = \text{Ker}([E_i, z]^{p-1} \text{ on } \tilde{P}_{i+1}) = 1$ . Since  $F_i \leq [E_i, z]^{p-1}$ , we have  $\text{Ker}(F_i \text{ on } \tilde{E}_{i+1}) = 1$ , that is  $D_i = F_i$ .

We first consider the case  $i = 2$ . Then  $p_2 = p$  and so  $p_1 \neq p$ . Since  $E_1 = P_1$  and  $[E_1, z] \neq 1$ , we see that  $\tilde{E}_2 \neq 0$ . Applying Theorem 2 to the action of  $E_1 < z >$  on each irreducible  $E_1 < z >$ -component of  $\tilde{E}_2$ , we get  $[\tilde{E}_2, z]^{p-1} \neq 0$ . This yields that  $[E_2, z]^{p-1} \neq 1$  and so  $F_2 = C_{[E_2, z]^{p-1}}(z) \neq 1$ . As  $F_1 = 1$ , this completes the proof of Claim 2 when  $i = 2$ .

We next assume that  $i > 2$ . Now  $p_{i-1} \neq p$  and  $F_{i-1} = C_{E_{i-1}}(z)$ . Since  $D_{i-1} \neq 1$ ,  $F_{i-1} \neq 1$  and  $\tilde{E}_i \neq 0$ . We apply Theorem 2 to the action of  $E_{i-1} < z >$  on each irreducible  $E_{i-1} < z >$ -component of  $\tilde{E}_i$  to get  $[\tilde{E}_i, z]^{p-1} \neq 0$  and  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } [\tilde{E}_i, z]^{p-1})$ . This gives that  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } [[\tilde{E}_i, z]^{p-1}, F_{i-1}])$  as  $[\tilde{E}_i, z]^{p-1} = [[\tilde{E}_i, z]^{p-1}, F_{i-1}] \oplus C_{[\tilde{E}_i, z]^{p-1}}(F_{i-1})$ . Now  $(F_{i-1} \text{ on } \tilde{E}_i) \equiv_w (F_{i-1} \text{ on } \tilde{F}_i)$  holds, because  $[\phi(E_i), F_{i-1}] = 1$ . This finishes the proof of Claim 2.  $\square$

Now we can prove the main result of this paper:

**Theorem A:** Let  $A$  be a finite abelian group with square free odd exponent not divisible by 3. Suppose that it acts fixed point freely on a finite group  $G$  of odd order. Then  $f(G) \leq \ell(A)$ .

**Proof:** Set  $f = f(G)$ . By Lemmas 8.1 and 8.2 in [2], there is an  $A$ -Fitting chain of

length  $f$  in  $G$ . Since  $A$  is nilpotent, it is a Carter subgroup of any semidirect product of it with a section of  $G$ . Thus  $A$  acts fixed point freely on any section of this chain.

Hence once the following assertion referring only to  $A$ -Fitting chains is proved, the theorem will follow immediately.

*Let  $A$  be finite abelian group of square free odd exponent not divisible by 3, and let  $P_1, \dots, P_t$  be an  $A$ -Fitting chain of a finite solvable group  $G$  such that  $P_i$  has odd order and  $A$  acts fixed point freely on  $P_i$  for each  $i = 1, \dots, t$ . Then  $t \leq \ell(A)$ .*

We shall use induction on  $t$ . We may assume that  $P_1$  is an irreducible  $A$ -module. As  $A$  acts fixed point freely on  $P_1$ , there exists  $z \in A$  of prime order  $p$  such that  $[P_1, z] \neq 1$ . Then  $[P_1, z] = P_1$  and so  $p_1 \neq p$ . Also  $p \geq 5$ . Theorem 3 applied to the chain  $P_1, \dots, P_t$  gives us an  $A$ -Fitting chain  $D_2, \dots, D_t$  such that  $z$  centralizes each  $D_i$ , for  $i = 2, \dots, t$ . Hence  $D_2, \dots, D_t$  is an  $A/\langle z \rangle$ -Fitting chain on each of its sections  $A/\langle z \rangle$  acts fixed point freely. By induction, it follows that  $t - 1 \leq \ell(A) - 1$ . Then  $t \leq \ell(A)$ , as desired.

**Lemma 4:** Let a finite group  $A$  act on a Fitting chain  $P_1, \dots, P_t$  where each  $P_i$  has odd order, in such a way that  $A$  centralizes no nontrivial section of any  $P_j$ ,  $j = 1, \dots, t$ . Assume that  $A$  is supersolvable of odd order which is not divisible by 3. Then  $t \leq 2(2^{\ell(A)} - 1)$ .

**Proof:** We shall use induction on  $\ell(A)$ . If  $\ell(A) = 0$ , then  $A = 1$  and hence the theorem follows. Assume that  $\ell(A) > 0$ , and that the theorem is true for all smaller values of  $\ell(A)$ . Let  $B$  be a normal subgroup of  $A$  such that  $|B|$  is the largest prime dividing  $|A|$ . We may assume that  $B$  centralizes  $P_1, \dots, P_k$  where  $k \in \{1, \dots, t\}$ , and  $k$  is the largest such positive integer.

Now  $A/B$  acts on  $P_1, \dots, P_k$  and by induction we have  $k \leq 2(2^{\ell(A/B)} - 1)$ . Since  $t = k + (t - k)$ , we may assume that  $t > 2^{\ell(A)}$ . Then  $t - k$ , the length of  $P_{k+1}, \dots, P_t$ , is at least 3. Then by Theorem 3 applied to  $P_{k+1}, \dots, P_t$ , we get a chain  $D_{k+3}, \dots, D_t$  of sections such that each  $D_j$  is centralized by  $B$ .

Since  $A/B$  and  $D_{k+3}, \dots, D_t$  fulfill the hypothesis, we see that

$$t - (k+3) + 1 \leq 2(2^{\ell(A)-1} - 1) \text{ and so } t \leq k + 2^{\ell(A)} \leq 2(2^{\ell(A)-1} - 1) + 2^{\ell(A)} = 2(2^{\ell(A)} - 1),$$

as desired.  $\square$

**Theorem B:** Let  $H$  be a finite group of odd order which is not divisible by 3. Suppose that its Carter subgroups have a normal complement  $G$ . If  $C$  is a Carter subgroup of  $H$ , then  $f(G) \leq 2(2^{\ell(C)} - 1)$ .

**Proof:** Set  $f = f(G)$ . By Lemmas 8.1 and 8.2 in [2], there is a C-Fitting chain  $P_f, \dots, P_1$ . Since  $C$  is a Carter subgroup of  $H$  with  $G \cap C = 1$ , it centralizes no nontrivial section of  $G$ . By Lemma 4, we obtain that  $f \leq 2(2^{\ell(C)} - 1)$ .  $\square$

**Theorem C:** Let  $C$  be a Carter subgroup of  $G$  where  $G$  is a finite group of odd order which is not divisible by 3. Then  $f(G) \leq 4(2^{\ell(C)} - 1) - \ell(C)$ .

**Proof:** Set  $f = f(G)$ . We use induction on  $\ell(C)$ . If  $\ell(C) = 0$ , then  $C = 1$ ,  $G = 1$  and so the theorem follows. Assume that  $\ell(C) > 0$  and that the theorem is true for all smaller values of  $\ell$ . Fix a Carter subgroup  $C$  of  $G$ . There is an integer  $k \geq 0$  such that  $F_k(G) \cap C = 1$  and  $F_{k+1}(G) \cap C \neq 1$ . Put  $\bar{G} = G/F_{k+1}(G)$ . Since  $\bar{C}$  is a Carter subgroup of  $\bar{G}$  and  $F_{k+1}(G) \cap C \neq 1$ ,  $\ell(\bar{C}) < \ell(C)$ . So by induction

$$f(\bar{G}) = f - k - 1 \leq (2^{\ell(\bar{C})} - 1) - \ell(\bar{C}) \leq 4(2^{\ell(C)-1} - 1) - (\ell(C) - 1)$$

Now  $C$  is a Carter subgroup of  $K = CF_k(G)$  and  $F_k(G)$  is a normal complement to each Carter subgroup of  $K$ . Thus  $k = f(F_k(G)) \leq 2(2^{\ell(C)} - 1)$  by Theorem B.

It follows that

$$f = 1 + k + (f - k - 1) \leq 1 + 2(2^{\ell(C)} - 1) + 4(2^{\ell(C)-1} - 1) - (\ell(C) - 1) = 4(2^{\ell(C)} - 1) - \ell(C).$$

### Acknowledgement

We are very grateful to Prof. Alexandre Turull for devoting his valuable time and effort to an earlier version of the article. His attentive reading and valuable suggestions helped us to improve the article in readability and gave a chance to correct a mistake in one of the proofs.

### References

- [1.] S.D.Bell and B.Hartley, A note on fixed point free actions of finite groups, Quart. J. Math. Oxford Ser. (2) 41 (1990), 127-130.

- [2.] E.C.Dade, Carter subgroups and Fitting heights of finite solvable groups, Illinois J. Math. 13 (1969), 449-514.
- [3.] G.Ercan and İ. Gülođlu, On finite groups admitting a fixed point free automorphism of order  $pqr$ , J. Group Theory 7 (2004), no. 4, 437-446.
- [4.] B.Huppert and N.Blackburn, Finite groups, vol.2 (Springer-Verlag, 1981).
- [5.] C.Kei-Nah, Finite groups admitting automorphisms of order  $pq$ , Proc. Edinburgh Math. Soc. (2) 30 (1987), 51-56.
- [6.] A.Turull, Fitting height of groups and of fixed points, J. Algebra 86 (1984), 555-566.
- [7.] A.Turull, Character theory and length problems, Finite and locally finite groups (İstanbul, 1994), 377-400.